

Chitosan as a growth stimulator in orchid tissue culture

Khin Lay Nge^a, Nitar Nwe^a, Suwalee Chandkrachang^{a,b}, Willem F. Stevens^{a,b,*}

^a *Bioprocess Technology, Food Engineering and Bioprocess Technology, Asian Institute of Technology, P.O. Box 4, Klong Luang, Pathumthani 12120, Bangkok, Thailand*

^b *Center for Chitin and Chitosan Biomaterials, Chulalongkorn University, Bangkok, Thailand*

Received 22 October 2005; received in revised form 8 February 2006; accepted 10 February 2006

Available online 10 March 2006

Abstract

The effect of shrimp and fungal chitosan on the growth and development of orchid plant meristemic tissue in culture was investigated in liquid and on solid medium. The growth of meristem explants into protocorm-like bodies in liquid medium was accelerated up to 15 times in the presence of chitosan oligomer, the optimal concentration being 15 ppm. The 1 kDa shrimp oligomer was slightly more effective compared to 10 kDa shrimp chitosan and four times more active compared to high molecular weight 100 kDa shrimp chitosan. The 10 kDa fungal chitosan was more effective compared with 1 kDa oligomer. The development of orchid protocorm into differentiated orchid tissue with primary shoots and roots was studied on solid agar medium. The optimal effect, the generation of 5–7 plantlets in 12 weeks was observed in the presence of 20 ppm using either 10 kDa fungal or 1 kDa oligomer shrimp chitosan. The data are consistent with preliminary results from field experiments and confirm unequivocally that a minor amount of chitosan has a profound effect on the growth and development of orchid plant tissue.

© 2006 Elsevier Ireland Ltd. All rights reserved.

Keywords: Fungal chitosan; Meristematic bud; Orchids; Protocorm; Shrimp chitosan

1. Introduction

Chitin is a natural polysaccharide, which consists of a copolymer of *N*-acetyl- D -glucosamine and D -glucosamine residues, linked by β -1,4 glycosidic bonds. It is present in a variety of species: in shells of crustaceans, in cuticles of insects and in the cell wall of fungi and some algae. The deacetylated form of chitin is chitosan. The chitosan biopolymer has applications in waste water treatment, pulp and paper, in medical and cosmetic products, biotechnology, food and feed and membranes [1]. In agriculture, chitosan has been used in seed, leaf, fruit and vegetable coating [2], as fertilizer and in controlled agrochemical release [3], to increase plant product [4–6], to stimulate the immunity of plants [7], to protect plants against microorganisms [8–10] and to stimulate plant growth. In the latter studies, a positive effect of chitosan was observed on the growth of roots, shoots and leaves of various plants including *Gerbera* [4] and of several crop plants [11]. Orchid roots sprayed with a very diluted chitosan solution show

stimulation of growth, renewed flower production and enhanced resistance against fungi and virus [5].

Among the commercially important orchids, *Dendrobium* accounts for about 80% of the total of micropropagated tropical orchids [12]. *Dendrobium* plants are used for the production of orchid flowers [13] and in traditional Chinese medicine [14]. *Dendrobium* is micropropagated in tissue culture by protocorm-like bodies (PLB) but the growth is very slow [15]. In order to stimulate efficient micropropagation of PLB, much effort has been directed to modify the culture media, mainly by the inclusion of plant growth regulators [16,17] such as *N*⁶-benzylaminopurine and α -naphthaleneacetic acid [18], benzyladenine, gibberellic acid and 3-indoleacetic acid [19], various polyamines [20], *N*⁶-benzyladenine and thidiazuron [21,22].

Plant tissue culture is particular suitable to study the effect of chitosan on plants that grow and multiply slowly, like orchids. In the work presented here, the formation of protocorm-like bodies in meristem buds and the growth and differentiation of orchid protocorm have been studied under the controlled conditions of tissue culture in the absence and presence of chitosan. Various chitosan preparations, both of crustacean and fungal origin have been compared in various concentrations.

* Corresponding author. Tel.: +66 2 969 3032; fax: +66 2 969 3033.

E-mail address: wfstevens@gmail.com (W.F. Stevens).

The formation of differentiated orchid tissue has been studied in liquid and on solid media. Precise data on the effectiveness of chitosan as growth stimulator are presented.

2. Materials and methods

2.1. Preparation of meristematic auxillary buds and protocorm-like bodies for explants

Young *Dendrobium phalaenopsis* orchid plants were obtained from Orchid Garden, Ratchaburi, Thailand. Shoot segments with one to two meristematic auxillary buds (6–15 mm) were excised. These shoot pieces were washed with soap and tap water three times and immersed in antiseptic solution (5.25% sodium hypochlorite with a few drops of soap) to disinfect the surface. After that the shoot segments were rinsed several times in sterilized distilled water to remove the antiseptic solution and non-attached tissue. Meristematic auxillary buds were excised from these shoot segments and cultured in liquid modified VW medium [23]. *Dendrobium* orchid protocorms three months of age were obtained from the Plant Tissue Culture Laboratory, Chatuchak Weekend Market, Bangkok, Thailand.

2.2. Culture medium and conditions

The VW medium [23] contained 20 g/l sucrose, 150 ml/l coconut water, 0.525 g/l KNO_3 , 0.25 g/l KH_2PO_4 , 0.5 g/l $(\text{NH}_4)_2\text{SO}_4$, 0.007 g/l $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.025 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.028 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ and 0.037 g/l $\text{Na}_2\text{-EDTA}$. Coconut water was obtained from local market, Bangkok, Thailand. The media were supplemented with various amounts of chitosan, either chitosan from shrimp shell with a molecular weight of 100 or 10 kDa [24]. Oligomer 1 kDa chitosan was obtained from Kittolife Co. Ltd., Korea. The 10 kDa chitosan was isolated from fungal cell wall [25]. The pH of the medium was adjusted to 4.9 for liquid medium. Solid medium was prepared by adding agar (12 g/l supplied by Sigma) to the complete liquid medium. Liquid medium (50 ml) in the 250 ml Erlenmeyer flasks and 10 ml of solid agar medium in the culture tubes (150 mm × 25 mm) were autoclaved for 20 min at 121 °C.

Meristematic auxillary buds were used as explants in the liquid medium. Protocorm-like bodies (PLB) (0.03–0.04 g fresh weight) were used as explant in liquid and agar medium. Liquid cultures were maintained at 27 °C and 100 rpm. Solid cultures were kept at 27 °C and provided photoperiods of 16 h per day using cool white fluorescent tubes (TLD 18 W/54 φ 4A, Daylight 1030 lm, 57 lm/W, Philips, Thailand).

2.3. Data collection and analysis

Visual observations were made every day. Total fresh weight of protocorm-like bodies was recorded at 0 and after 3, 6, 8 and 12 weeks. The number of plantlets generated by the protocorm-like bodies maintained on agar medium were counted after 12 weeks of cultivation. Three replicates were

carried out in each experiment. All data were analyzed by SPSS software, version 7.2 using one-way ANOVA analysis. All photographs are representative for the growth condition as described.

3. Results and discussion

3.1. Effect of shrimp chitosan on orchid meristematic bud growth in liquid medium

Meristematic buds of mature orchid plants form initially a protocorm-like body (PLB) when cultivated in VW liquid medium. The growth of meristematic tissue is dependent on size of the inoculum and on the conditions of growth. PLB were grown in liquid media supplemented with 15 ppm of shrimp chitosan of different molecular weights (100, 10 kDa and oligomer 1 kDa). After three weeks, orchid meristem buds cultures supplemented with oligomer chitosan showed more cell propagation than the cultures supplemented with 10 and 100 kDa chitosan ($p < 0.05$). The cell mass in the former culture medium increased to about 15 times after six weeks cultivation. High molecular weight chitosan (100 kDa) had no stimulating effect (Fig. 1).

The effect of chitosan concentration on the growth of PLB's from meristematic buds was investigated in liquid medium supplemented with 0, 5, 10, 15, 20 and 25 ppm of oligomeric chitosan. The medium supplemented with 10 and 15 ppm produced PLB larger in size in comparison to the other treatments. Maximum fresh weight of PLB's was obtained in culture media supplemented with 15 ppm oligomer chitosan (Fig. 2). Under this condition the excised meristem tissue initially increased in size and changed into small round bodies during the first three weeks, after that juvenile leaves appeared after five weeks (Fig. 3). In media with 5, 20 and 25 ppm oligomer chitosan PLB, increase in size was not observed after six weeks of cultivation.




Fig. 1. Effect of shrimp chitosan with different molecular weights. Orchid protocorm-like bodies (PLB's) were cultivated during 0, 3 and 6 weeks on VW liquid medium supplemented with shrimp chitosan with molecular weights of 1 kDa (oligo-chitosan), and of 10 and 100 kDa. Standard deviation was calculated from three independent experiments by one-way ANOVA.




Fig. 2. Effect of different concentrations of oligomer chitosan on the weight of PLB's after 0, 21 and 42 days in culture in VW liquid medium. Standard deviation was calculated from three independent experiments by one-way ANOVA.

3.2. Effect of chitosan from shrimp and fungal origin on the growth of PLB in liquid medium

The effect of chitosan preparations from shrimp and fungal origin on PLB was compared using 1 kDa shrimp chitosan, 10 kDa shrimp chitosan and 10 kDa fungal chitosan in liquid medium in concentrations of 0–5 ppm. The increase in fresh weight of PLB's was registered. The results are shown in Fig. 4. The protocorms (original weight 0.03–0.04 g) in the media supplemented with oligomer and fungus chitosan at three concentration viz., 10, 15 and 20-ppm produced higher weight PLB's in comparison to control media without chitosan ($p < 0.05$). After six weeks, media with 15 ppm fungal chitosan showed a stimulated 30-fold increase in fresh weight in comparison with PLB weight at the start of the experiment. Over the same period of time in control cultures a non-stimulated increase in weight of 5–6-fold was observed. This non-stimulated increase was also observed in cultures of 1 kDa oligo chitosan and 10 kDa fungal chitosan in concentrations of 5 and 25 ppm and in all cultures with 10 kDa chitosan.

In the media supplemented with 10 kDa shrimp chitosan a small brown precipitate was observed after two days of incubation. However the color and transparency of the liquid




Fig. 4. Effect of various concentrations of shrimp and fungal chitosan on the propagation of total fresh weight of PLB's after six weeks cultivation in liquid VW medium. Shrimp chitin 10 kDa, shrimp oligomer 1 kDa, fungal chitosan 10 kDa. Standard deviation was calculated from three independent experiments by one-way ANOVA.

media did not change during subsequent cultivation during six weeks. Also in the fungal chitosan supplemented media a very small amount of dark precipitate was formed after two days but the solutions remained clear thereafter. After eight weeks investigation, 15 ppm of fungus chitosan seemed the most effective in propagation of PLB's as measured by the increase in their fresh weight (Fig. 5). Therefore fungus chitosan is an attractive candidate to be used as growth stimulator in VW liquid medium for orchid plant tissue culture.

3.3. Effect of various chitosan on differentiation on agar medium

In liquid media, orchid protocorm propagated only by increasing its total fresh weight. However, new plantlets of orchid can be obtained by cultivation of orchid protocorm in static cultures on the surface of solid agar media. In this system, the effect of 1 and 10 kDa shrimp chitosan and 10 kDa fungal chitosan were compared in the range of 0–25 ppm.

An important criterion in plantlet development is the formation of fresh-green differentiated tissue. Green color tissue was observed in the protocorms after two to three weeks on the medium supplied with chitosan. The green differentiated




Fig. 3. Effect of shrimp oligomer chitosan (15 ppm) on the development of orchid protocorm from the meristematic bud after six weeks of cultivation in VW medium.




Fig. 5. Effect of fungal chitosan (15 ppm) on the propagation of PLB tissue after eight weeks cultivation in VW medium.



Fig. 6. Effect of various chitosan preparations (20 ppm) during 12 weeks cultivation on VW agar medium on the generation of plantlets by PLB explant of *Dendrobium phalaenopsis* in VW liquid medium.

tissue on the protocorm on the control media was observed only after six weeks of cultivation.

Small plantlets started to develop from the green protocorm after five to six weeks of cultivation. After a cultivation period of 12 weeks, the number of plantlets on the media supplied with 10 kDa fungus chitosan was the highest compared to other chitosan preparations (Fig. 6). Precise data for the effect of three chitosan preparations in five concentrations after 12 weeks of cultivation are presented in Table 1. Five to seven plantlets were generated from the green protocorm on the media supplied with 20 ppm of 10 kDa fungal or 1 kDa oligomer chitosan. Protocorm on the media supplied with 5 and 10-ppm fungal chitosan did not form plantlets during 12 weeks cultivation. Representative pictures of this cultivation experiment are shown in Fig. 7. A similar influence of low molecular weight chitosan in vitro regeneration of somatic embryos was observed in regeneration of callus in carrot and root and shoot regeneration in strawberry [26]. The best solid medium for the cultivation of

orchid tissue culture is the medium supplied with 20 ppm of 10 kDa fungal or 1 kDa oligomer chitosan.

3.4. Comparison with data from field experiments

The data in tissue culture presented here support earlier preliminary observations made in the field, that spraying of low molecular weight chitosan in the dosage of 15–20 ppm to orchid plant results in a considerable positive effect on plant growth [4–6,11]. In addition the chitosan acts as antifungal compound. The data confirm unequivocally that a minor amount of chitosan has profound effects on the growth and development of orchid plant tissue [8–10]. Chitosan has a unique combination of attractive characteristics: it stimulates plant growth, it provides plant protection, it is environmental friendly being of biological origin and is easy biodegradable by soil microorganisms. The unique combination of these properties makes chitosan to a very useful biocontrol agent with a large perspective for horticulture in general and orchid culture in particular.

Table 1

Number of plantlets generated by orchid protocorm grown on VW agar media supplemented with different concentrations of shrimp chitosan, chitosan oligomer and fungal chitosan after 12 week in cultivation

Chitosan	5 ppm	10 ppm	15 ppm	20 ppm	25 ppm	Control (0 ppm)
10 kDa shrimp	0 ± 0	0 ± 0	1.7 ± 1.5	4.7 ± 0.6	1.3 ± 0.6	1.3 ± 1.2
1 kDa oligomer	1.7 ± 0.6	1.7 ± 1.5	3.0 ± 2.7	6.3 ± 0.6	2.7 ± 2.3	1.3 ± 1.2
10 kDa fungus	0.3 ± 0.6	0.7 ± 0.6	3.3 ± 1.2	6.7 ± 0.6	4.0 ± 3.5	1.3 ± 1.2

All data are the average of three replicate independent experiments and the standard deviation was calculated using one-way ANOVA.




Fig. 7. Effect of various concentrations of fungal chitosan (10 kDa) on the production of plantlets by the fractionated PLB's explant of *Dendrobium phalaenopsis* cultivated on VW agar media. Photographs taken at 12 weeks after inoculation.

4. Conclusion

Different chitosan preparations were investigated for their effect on orchid meristem culture and orchid protocorm propagation in liquid and on agar media. According to the results obtained, chitosan has the ability to stimulate the induction of the differentiation of orchid plant tissue. Fungal chitosan and shrimp oligomer chitosan of 15 and 20 ppm are most effective on protocorm propagation in liquid and on agar orchid plant tissue culture media, respectively.

Acknowledgements

The authors thank the late Prince de Lignac, Prospect Burma and Open Society Institute for providing funds, Mr. Suwitjai Sangtein (Orchid Garden, Ratchaburi, Thailand) for providing the young orchid plants and Dr. Oradee Sahavacharin for teaching the tissue culture technique.

References

- [1] S. Hirano, Applications of chitin and chitosan in the ecological and environment fields, in: M.F.A. Goosen (Ed.), Application of Chitin and Chitosan, Technomic Publishing Company, 1997, pp. 31–54, ISBN 1-56676-449-1.
- [2] F. Devlieghere, A. Vermeulen, J. Debevere, Chitosan: antimicrobial activity, interactions with food components and applicability as a coating on fruit and vegetables, Food Microbiol. (2004) 703–714.
- [3] M. Sukwattanasinitt, A. Klaikherd, K. Skulnee, S. Aiba, Chitosan as a releasing device for 2,4-D herbicide, in: T. Urugami, K. Kurita, T. Fukamizo (Eds.), Chitin and Chitosan, Chitin and Chitosan in Life Science, Yamaguchi, 2001, pp. 142–143, ISBN 4-906464-43-0.
- [4] P. Wanichpongpan, K. Suriyachan, S. Chandkrachang, Effects of chitosan on the growth of Gerbera flower plant (*Gerbera jamesonii*), in: T. Urugami, K. Kurita, T. Fukamizo (Eds.), Chitin and Chitosan in Life Science, Yamaguchi, 2001, pp. 198–201, ISBN 4-906464-43-0.
- [5] S. Chandkrachang, The applications of chitin and chitosan in agriculture in Thailand, in: K. Suchiva, S. Chandkrachang, P. Methacanon, M.G. Peter (Eds.), Advances in Chitin Science, vol. 5, Bangkok, 2002, pp. 458–462, ISBN 974-229-412-7.
- [6] N. Nwe, S. Chandkrachang, W.F. Stevens, Application of chitosan in Myanmar's agriculture sector, in: Proceedings of the Sixth Asia Pacific Chitin and Chitosan Symposium, May 23–26, The National University of Singapore, Singapore, 2004.
- [7] L.A. Hadwiger, S.J. Klosterman, J.J. Choi, The mode of action of chitosan and its oligomers in inducing plant promoters and developing disease resistance in plants, in: K. Suchiva, S. Chandkrachang, P. Methacanon, M.G. Peter (Eds.), Advances in Chitin Science, vol. 5, Bangkok, 2002, pp. 452–457, ISBN 974-229-412-7.
- [8] H. Pospieszny, S. Chirkov, J. Atabekov, Induction of antiviral resistance in plants by chitosan, Plant Sci. 79 (1991) 63–68.
- [9] H. Struszczyk, H. Pospieszny, New applications of chitosan and its derivatives in plant protection, in: M.F.A. Goosen (Ed.), Applications of Chitin and Chitosan, Technomic Publishing Co., Inc., USA, 1997, pp. 171–184, ISBN 1-56676-449-1.
- [10] S. Bautista-Baños, M. Hernandez-Lopez, E. Bosquez-Molina, C.L. Wilson, Effects of chitosan and plant extracts on growth of *Colletotrichum gloeosporioides*, anthracnose levels and quality of papaya fruit, Crop Protect. 22 (2003) 1087–1092.
- [11] H. Chibu, H. Shibayama, Effects of chitosan applications on the growth of several crops, in: T. Urugami, K. Kurita, T. Fukamizo (Eds.), Chitin and Chitosan in Life Science, Yamaguchi, 2001, pp. 235–239, ISBN 4-906464-43-0.
- [12] P.C. Debergh, R.H. Zimmerman, Micropropagation: Technology and Application, Kluwer Academic Publishers, The Netherlands, 1991.
- [13] S. Ketsa, Y. Piyasaengthong, S. Prathuangwong, Mode of action of AgNO₃ in maximizing vase life of *Dendrobium 'Pompadour'* flowers, Postharvest Biol. Technol. (1995) 109–117.
- [14] C. Fan, W. Wang, Y. Wang, Q. Guowei, W. Zhao, Chemical constituents from *Dendrobium densiflorum*, Phytochemistry (2001) 1255–1258.
- [15] A.D. Hawkes, Encyclopedia of Cultured Orchids, Fabre and Fabre Limited, London, 1970.
- [16] L. Prakash, C.L. Lee, C.S. Loh, C.J. Goh, in: A.S. Islam (Ed.), In Vitro Propagation of Commercial Orchids: An Assessment of Current Methodologies and Development of a Novel Approach-Thin Cross-Section Culture, Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, 1996 pp. 42–49.

- [17] N.R. Nayak, S. Sahoo, S. Patnaik, S.P. Rath, Establishment of thin cross section (TCS) culture method for rapid micropropagation of *Cymbidium aloifolium* (L.) Sw. and *Dendrobium nobile* Lindl. (Orchidaceae), *Scientia Hort.* 94 (2002) 107–116.
- [18] J. Roy, N. Banerjee, Induction of callus and plant regeneration from shoot-tip explants of *Dendrobium fimbriatum* Lindl. Var. *oculatum* Hk. F., *Scientia Hort.* 97 (2003) 333–340.
- [19] C.J. Goh, A.L. Yang, Effects of growth regulators and decapitation on flowering of *Dendrobium* orchid hybrids, *Plant Sci., Lett.* (1978) 287–292.
- [20] G.V.S. Saiprasad, P. Raghuvver, S. Khetarpal, R. Chandra, Effect of various polyamines on production of protocorm-like bodies in orchid *Dendrobium* 'Sonia', *Scientia Hort.* (2004) 161–168.
- [21] N.R. Nayak, S.P. Rath, S. Patnaik, In vitro propagation of three epiphytic orchids, *Cymbidium aloifolium* (L.) Sw., *Dendrobium aphyllum* (Roxb.) Fisch. and *Dendrobium moschatum* (Buch-Ham) Sw. through thidiazuron-induced high frequency shoot proliferation., *Scientia Hort.* (1997) 243–250.
- [22] R.B. Malabadi, G.S. Mulgund, N. Kallappa, Micropropagation of *Dendrobium nobile* from shoot tip sections, *J. Plant Physiol.* 162 (2005) 473–478.
- [23] E.F. Vacin, F.W. Went, Some pH changes in nutrient solutions, *Bot. Gaz.* 110 (1949) 605–617.
- [24] W.F. Stevens, Production of chitin and chitosan: refinement and sustainability of chemical and biological processing, in: T. Uragami, K. Kurita, T. Fukamizo (Eds.), *Chitin and Chitosan in Life Science*, Yamaguchi, 2001, pp. 293–300, ISBN 4-906464-43-0.
- [25] N. Nwe, W.F. Stevens, Production of fungal chitosan by solid substrate fermentation followed by enzymatic extraction, *Biotechnol. Lett.* 24 (2002) 131–134.
- [26] L.Q. Luan, V.T.T. Ha, L. Hai, N.Q. Hien, N. Nagasawa, F. Yoshii, T. Kume, Study on the biological effect of irradiated chitosan on plant in tissue culture, in: K. Suchiva, S. Chandkrachang, P. Methacanon, M.G. Peter (Eds.), *Advances in Chitin Science*, vol. 5, Bangkok, 2002, pp. 468–474, ISBN 974-229-412-7.